Scientific article

DOI: 10.18287/2541-7525-2020-26-4-15-24

Submitted: 14.10.2020

Revised: 16.11.2020

Accepted: 25.11.2020

S. V. Astashkin

Samara National Research University, Samara, Russian Federation
 E-mail: astash56@mail.ru. ORCID: <https://orcid.org/0000-0002-8239-5661>

SYMMETRIC FINITE REPRESENTABILITY OF ℓ^p IN ORLICZ SPACES¹

ABSTRACT

It is well known that a Banach space need not contain any subspace isomorphic to a space ℓ^p ($1 \leq p < \infty$) or c^0 (it was shown by Tsirel'son in 1974). At the same time, by the famous Krivine's theorem, every Banach space X always contains at least one of these spaces *locally*, i.e., there exist finite-dimensional subspaces of X of arbitrarily large dimension n which are isomorphic (uniformly) to ℓ_p^n for some $1 \leq p < \infty$ or c_0^n . In this case one says that ℓ^p (resp. c^0) is finitely representable in X . The main purpose of this paper is to give a characterization (with a complete proof) of the set of p such that ℓ^p is *symmetrically* finitely representable in a separable Orlicz space.

Key words: ℓ^p -space; finite representability of ℓ^p -spaces; symmetric finite representability of ℓ^p -spaces; Orlicz function space; Orlicz sequence space; Matuszewska-Orlicz indices.

Citation. Astashkin S.V. Symmetric finite representability of ℓ^p in Orlicz spaces. *Vestnik Samarskogo universiteta. Estestvennonauchnaia seriia* = *Vestnik of Samara University. Natural Science Series*, 2020, vol. 26, no. 4, pp. 15–24. DOI: <http://doi.org/10.18287/2541-7525-2020-26-4-15-24>.

Information about the conflict of interests: author and reviewers declare no conflict of interests.

© Astashkin S.V., 2020

Astashkin Sergey Vladimirovich — Doctor of Physical and Mathematical Sciences, professor, head of the Department of Functional Analysis and Function Theory, Samara National Research University, 34, Moskovskoye shosse, 443086, Russian Federation.

Introduction

While a Banach space X need not contain any subspace isomorphic to a space ℓ^p ($1 \leq p < \infty$) or c^0 (as was shown by Tsirel'son in [1]), it will always contain at least one of these spaces *locally*. This means that there exist finite-dimensional subsets of X of arbitrarily large dimension n which are isomorphic (uniformly) to ℓ_p^n for some $1 \leq p < \infty$ or c_0^n . This fact is the content of the famous result proved by Krivine in [2] (see also [3]). To state it we need some definitions.

Suppose X is a Banach space, $1 \leq p \leq \infty$, and $\{z_i\}_{i=1}^\infty$ is a bounded sequence in X . The space ℓ^p is said to be *block finitely representable in* $\{z_i\}_{i=1}^\infty$ if for every $n \in \mathbb{N}$ and $\varepsilon > 0$ there exist $0 = m_0 < m_1 < \dots < m_n$ and $\alpha_i \in \mathbb{R}$ such that the vectors $u_k = \sum_{i=m_{k-1}+1}^{m_k} \alpha_i z_i$, $k = 1, 2, \dots, n$, satisfy the inequality

$$(1 + \varepsilon)^{-1} \|a\|_p \leq \left\| \sum_{k=1}^n a_k u_k \right\|_X \leq (1 + \varepsilon) \|a\|_p$$

for arbitrary $a = (a_k)_{k=1}^n \in \mathbb{R}^n$. In what follows,

$$\|a\|_p := \left(\sum_{k=1}^n |a_k|^p \right)^{1/p} \text{ if } p < \infty, \text{ and } \|a\|_\infty := \max_{k=1,2,\dots,n} |a_k|$$

The space ℓ^p , $1 \leq p \leq \infty$, is said to be *finitely representable* in X if for every $n \in \mathbb{N}$ and $\varepsilon > 0$ there exist $x_1, x_2, \dots, x_n \in X$ such that for any $a = (a_k)_{k=1}^n \in \mathbb{R}^n$

$$(1 + \varepsilon)^{-1} \|a\|_p \leq \left\| \sum_{k=1}^n a_k x_k \right\|_X \leq (1 + \varepsilon) \|a\|_p$$

¹The work was completed as a part of the implementation of the development program of the Scientific and Educational Mathematical Center the Volga Federal District, agreement no. 075-02-2021-1393.

(alternatively, in the case $p = \infty$, one might say that c^0 is finitely representable in X).

Clearly, if ℓ^p is block finitely representable in some sequence $\{z_i\}_{i=1}^\infty \subset X$, then ℓ^p is finitely representable in X . Therefore, the following famous result proved by Krivine in [2] (see also [3] and [4, Theorem 11.3.9]) implies the finite representability of ℓ^p for some $1 \leq p \leq \infty$ in any Banach space.

Theorem (Krivine)

Let $\{z_i\}_{i=1}^\infty$ be an arbitrary normalized sequence in a Banach space X such that the vectors z_i do not form a relatively compact set. Then ℓ^p is block finitely representable in $\{z_i\}_{i=1}^\infty$ for some $p \in [1, \infty]$.

Here, we consider both Orlicz sequence and function spaces (see the next section for the definition) and in the separable case we give a characterization of the set of p such that ℓ^p is *symmetrically finitely representable* in such a space. To introduce the notion of symmetric finite representability, we need some more definitions.

A sequence $y = (y_k)_{k=1}^\infty$ will be called a *copy* of a sequence $x = (x_k)_{k=1}^\infty$ if x and y have the same entries, that is, there is a permutation π of the set of positive integers such that $y_{\pi(k)} = x_k$ for all $k = 1, 2, \dots$

Given a measurable function $x(t)$ on $[0, 1]$, we set

$$n_x(\tau) := m(\{t \in [0, \alpha) : |x(t)| > \tau\}), \quad \tau > 0.$$

Here and in the sequel, m denotes the Lebesgue measure. Functions $x(t)$ and $y(t)$ are called *equimeasurable* if $n_x(\tau) = n_y(\tau)$ for each $\tau > 0$.

Let X be a symmetric sequence space (see e.g. [5]), $1 \leq p \leq \infty$. We say that ℓ^p is *symmetrically finitely representable* in X if for every $n \in \mathbb{N}$ and each $\varepsilon > 0$ there exists an element $x_0 \in X$ such that for its disjoint copies x_k , $k = 1, 2, \dots, n$, and for every $(a_k)_{k=1}^n \in \mathbb{R}^n$ we have

$$(1 + \varepsilon)^{-1} \|a\|_p \leq \left\| \sum_{k=1}^n a_k x_k \right\|_X \leq (1 + \varepsilon) \|a\|_p$$

Similar notion will be defined also in the function case. Let X be a symmetric function space on $[0, 1]$ [5]. The space ℓ^p is *symmetrically finitely representable* in X if for every $n \in \mathbb{N}$ and $\varepsilon > 0$ there exist equimeasurable and disjointly supported on $[0, 1]$ functions $u_i(t)$, $i = 1, 2, \dots, n$, such that for all $(a_k)_{k=1}^n \in \mathbb{R}^n$

$$(1 - \varepsilon) \|a\|_p \leq \left\| \sum_{i=1}^n a_i u_i \right\|_X \leq (1 + \varepsilon) \|a\|_p$$

The set of all $p \in [1, \infty]$ such that ℓ^p is symmetrically finitely representable in X (in both sequence and function cases) we will denote by $\mathcal{F}(X)$.

From the definition² of the Matuszewska-Orlicz indices α_N^0 and β_N^0 (resp. α_N^∞ and β_N^∞) of an Orlicz sequence space ℓ_N (resp. an Orlicz function space L_N) it follows that $\mathcal{F}(X) \subset [\alpha_N^0, \beta_N^0]$ (resp. $\mathcal{F}(X) \subset [\alpha_N^\infty, \beta_N^\infty]$). The main purpose of this paper is to give a detailed proof of the opposite embedding for both Orlicz sequence and function spaces. To this end, following the idea mentioned in [6, p. 140–141] we will make use of the proof of Theorem 4.a.9 from [7].

Similar problems for Orlicz function spaces (and more generally symmetric spaces) on $(0, \infty)$ were considered in the paper [8].

1. Preliminaries

1.1. Orlicz sequence spaces

A detailed information related to Orlicz sequence and function spaces see in monographs [9–11].

The Orlicz sequence spaces are a natural generalization of the ℓ^p -spaces, $1 \leq p \leq \infty$, which equipped with the usual norms

$$\|a\|_{\ell^p} := \begin{cases} (\sum_{k=1}^\infty |a_k|^p)^{1/p}, & 1 \leq p < \infty \\ \sup_{k=1,2,\dots} |a_k|, & p = \infty. \end{cases}$$

Let N be an Orlicz function, that is, an increasing convex continuous function on $[0, \infty)$ such that $N(0) = 0$ and $\lim_{t \rightarrow \infty} N(t) = \infty$. The *Orlicz sequence space* ℓ_N consists of all sequences $a = (a_k)_{k=1}^\infty$, for which the following (Luxemburg) norm

$$\|a\|_{\ell_N} := \inf \left\{ u > 0 : \sum_{k=1}^\infty N\left(\frac{|a_k|}{u}\right) \leq 1 \right\}$$

²See the next section.

is finite. Without loss of generality, we will assume that $N(1) = 1$. In particular, if $N(t) = t^p$, we get the ℓ^p -space, $1 \leq p < \infty$.

Recall that the Matuszewska-Orlicz indices (at zero) α_N^0 and β_N^0 of an Orlicz function N are defined by

$$\alpha_N^0 := \sup \left\{ p : \sup_{x, y \leq 1} \frac{N(x)y^p}{N(xy)} < \infty \right\}, \quad \beta_N^0 := \inf \left\{ p : \inf_{x, y \leq 1} \frac{N(x)y^p}{N(xy)} > 0 \right\}.$$

It can be easily checked that $1 \leq \alpha_N^0 \leq \beta_N^0 \leq \infty$. It is well known also that an Orlicz sequence space ℓ_N is separable if and only if $\beta_N^0 < \infty$, or equivalently, if the function N satisfies the Δ_2 -condition at zero, i.e.,

$$\limsup_{u \rightarrow 0} \frac{N(2u)}{N(u)} < \infty.$$

The subset h_N of an Orlicz sequence space ℓ_N consists of all $(a_k)_{k=1}^\infty \in \ell_N$ such that

$$\sum_{k=1}^\infty N\left(\frac{|a_k|}{u}\right) < \infty \text{ for each } u > 0.$$

One can easily check (see also [7, Proposition 4.a.2]) that h_N is a separable closed subspace of ℓ_N and the canonical unit vectors $e_n = (e_n^i)$ such that $e_n^n = 1$ and $e_n^i = 0$ if $i \neq n$, $n = 1, 2, \dots$, form a symmetric basis of the space h_N . Recall that a basis $\{x_n\}_{n=1}^\infty$ of a Banach space X is said to be *symmetric* if there exists $C > 0$ such that for any permutation π of the set of positive integers and all $a_n \in \mathbb{R}$ we have

$$C^{-1} \left\| \sum_{n=1}^\infty a_n x_n \right\|_X \leq \left\| \sum_{n=1}^\infty a_n x_{\pi(n)} \right\|_X \leq C \left\| \sum_{n=1}^\infty a_n x_n \right\|_X.$$

Observe that the definition of an Orlicz sequence space ℓ_N is determined (up to equivalence of norms) by the behaviour of the function N near zero. More precisely, the following conditions are equivalent: 1) $\ell_N = \ell_M$ (with equivalence of norms); 2) the canonical vector bases of the spaces h_N и h_M are equivalent; 3) there are constants $C > 0$, $c > 0$ and $t_0 > 0$ such that for all $0 \leq t \leq t_0$ it holds

$$cN(C^{-1}t) \leq M(t) \leq c^{-1}N(Ct)$$

(see e.g. [7, Proposition 4.a.5] or [11, Theorem 3.4]). In particular, if N is a degenerate Orlicz function, i. e., for some $t_0 > 0$ we have $N(t) = 0$ if $0 \leq t \leq t_0$, then $\ell_N = \ell_\infty$ (with equivalence of norms).

Given Orlicz function N , we define the following subsets of the space $C[0, \frac{1}{2}]$:

$$E_{N,a}^0 := \overline{\left\{ \frac{N(xy)}{N(y)} : 0 < y < a \right\}}, \quad E_N^0 := \bigcap_{0 < a < 1} E_{N,a}^0$$

and

$$C_{N,a}^0 := \overline{\text{conv} E_{N,a}^0}, \quad C_N^0 := \bigcap_{0 < a < 1} C_{N,a}^0,$$

where $0 < a < 1$ and the closure is taken in the norm topology of $C[0, \frac{1}{2}]$. All these sets are non-void norm compact subsets of the space $C[0, \frac{1}{2}]$ [7, Lemma 4.a.6]. It is well known that they determine to a large extent the structure of disjoint sequences of Orlicz sequence spaces (see [7, § 4.a] and [12]). Moreover, if $1 \leq p < \infty$, then $t^p \in C_N^0$ if and only if $p \in [\alpha_N^0, \beta_N^0]$ [7, Theorem 4.a.9].

In the case when an Orlicz function N satisfies the Δ_2 -condition at zero, the sets $E_{N,a}^0$, E_N^0 , $C_{N,a}^0$ and C_N^0 can be considered as subsets of the space $C[0, 1]$ (see the remark after Lemma 4.a.6 in [7]).

1.2. Orlicz function spaces

Let N be an Orlicz function such that $N(1) = 1$. Denote by L_N the Orlicz space on $[0, 1]$ endowed with the Luxemburg norm

$$\|x\|_{L_N} := \inf \left\{ u > 0 : \int_0^1 N\left(\frac{|x(t)|}{u}\right) dt \leq 1 \right\}.$$

In particular, if $N(t) = t^p$, $1 \leq p < \infty$, we obtain the space $L_p = L_p[0, 1]$ with the usual norm.

The Matuszewska-Orlicz indices α_N^∞ and β_N^∞ (at infinity) of an Orlicz function N are defined by the formulae

$$\alpha_N^\infty = \sup \left\{ p : \sup_{x, y \geq 1} \frac{N(x)y^p}{N(xy)} < \infty \right\}, \quad \beta_N^\infty = \inf \left\{ p : \inf_{x, y \geq 1} \frac{N(x)y^p}{N(xy)} > 0 \right\}.$$

Again $1 \leq \alpha_N^\infty \leq \beta_N^\infty \leq \infty$. As in the case of sequence spaces, an Orlicz space L_N is separable if and only if $\beta_N^\infty < \infty$, or equivalently, if the function N satisfies the Δ_2 -condition at infinity, i.e.,

$$\limsup_{u \rightarrow \infty} \frac{N(2u)}{N(u)} < \infty.$$

In contrast to the sequence case, the definition of an Orlicz function space L_N on $[0, 1]$ is determined (up to equivalence of norms) by the behaviour of the function $N(t)$ for large values of t .

For every Orlicz function N we define the following subsets of the space $C[0, \frac{1}{2}]$:

$$E_{N,A}^\infty := \overline{\left\{ \frac{N(xy)}{N(y)} : y > A \right\}}, E_N^\infty = \bigcap_{A>0} E_{N,A}^\infty, \quad C_N^\infty := \overline{\text{conv} E_N^\infty},$$

where the closure is taken in the norm topology of $C[0, \frac{1}{2}]$. Again all these sets are non-void norm compact subsets of the space $C[0, \frac{1}{2}]$ and they determine largely the structure of disjoint sequences in Orlicz function spaces (see [12, Propositions 3 and 4]). Moreover, if $1 \leq p < \infty$, then $t^p \in C_N^\infty$ if and only if $p \in [\alpha_N^\infty, \beta_N^\infty]$ [12].

Finally, if an Orlicz function N satisfies the Δ_2 -condition at infinity, the sets $E_{N,A}^\infty$, E_N^∞ and C_N^∞ can be considered as subsets of the space $C[0, 1]$.

2. Symmetric finite representability of ℓ^p in Orlicz sequence spaces

Theorem 1

Let M be an Orlicz function satisfying the Δ_2 -condition at zero. Then ℓ^p is symmetrically finitely representable in the Orlicz sequence space ℓ_M if and only if $p \in [\alpha_M^0, \beta_M^0]$, i.e., $\mathcal{F}(\ell_M) = [\alpha_M^0, \beta_M^0]$.

Proof.

As was observed in §1, we always have $\mathcal{F}(\ell_M) \subset [\alpha_M^0, \beta_M^0]$. Therefore, it suffices to prove only the opposite embedding. In other words, we need to show that for every $p \in [\alpha_M^0, \beta_M^0]$, $m \in \mathbb{N}$ and each $\varepsilon > 0$ there exists an element $x_0 \in \ell_M$ such that for its disjoint copies x_k , $k = 1, 2, \dots, m$, and for every $c = (c_k)_{k=1}^m \in \mathbb{R}^n$ we have

$$(1 + \varepsilon)^{-1} \|c\|_p \leq \left\| \sum_{k=1}^m c_k x_k \right\|_{\ell_M} \leq (1 + \varepsilon) \|c\|_p. \quad (1)$$

According to the proof of Theorem 4.a.9 in [7] and a comment followed this proof on p. 144, $t^p \in C_M^0$ (see also §2.1). Since M satisfies the Δ_2 -condition at zero, the set C_M^0 may be considered as a subset of the space $C[0, 1]$ (see the remark after Lemma 4.a.6 in [7] or again §2.1). Therefore, since $C_M^0 := \bigcap_{0 < a < 1} C_{M,a}^0$, we conclude that $t^p \in C_{M,2^{-n}}^0$ for each $n \in \mathbb{N}$.

Note that the mapping

$$\lambda \mapsto M_\lambda(t) := M(\lambda t)/M(\lambda) \quad (2)$$

is continuous from $I_n := (0, 2^{-n}]$ into the subset $E_{M,2^{-n}}^0$ of $C[0, 1]$. Indeed, as it is well known (see e.g. [9, Theorem 1.1]),

$$M(t) = \int_0^t \rho(s) ds, \quad (3)$$

where ρ is a nondecreasing right-continuous function.

Therefore, for arbitrary $\lambda_2 > \lambda_1 > 0$ and all $0 \leq t \leq 1$ we have

$$\begin{aligned} |M_{\lambda_2}(t) - M_{\lambda_1}(t)| &= \frac{|M(\lambda_1)M(\lambda_2 t) - M(\lambda_2)M(\lambda_1 t)|}{M(\lambda_1)M(\lambda_2)} \\ &\leq \frac{1}{M(\lambda_2)} (M(\lambda_2 t) - M(\lambda_1 t) + M(\lambda_2) - M(\lambda_1)) \\ &\leq \frac{1}{M(\lambda_2)} \left(\int_{\lambda_1 t}^{\lambda_2 t} \rho(s) ds + \int_{\lambda_1}^{\lambda_2} \rho(s) ds \right) \\ &\leq \frac{2\rho(\lambda_2)}{M(\lambda_2)} (\lambda_2 - \lambda_1). \end{aligned}$$

Thus, mapping (2) may be extended uniquely to a map $\omega \mapsto M_\omega$ from the Stone-Ćech compactification βI_n of I_n onto the set $E_{M,2^{-n-1}}^0$. Since $t^p \in C_{M,2^{-n}}^0$ and the extreme points of $C_{M,2^{-n}}^0$ are contained in the

compact set $E_{M,2^{-n}}^0$, by the Krein-Milman theorem (see e.g. [13, Theorem 3.28]), there exists a probability measure μ_n on the set βI_n such that

$$t^p = \int_{\beta I_n} M_\omega(t) d\mu_n(\omega), \quad 0 \leq t \leq 1. \quad (4)$$

Let us show that

for some probability measure ν_n on I_n we have

$$\left| t^p - \int_0^{2^{-n}} M_\lambda(t) d\nu_n(\lambda) \right| < 2^{-n}, \quad 0 \leq t \leq 1. \quad (5)$$

First, the fact that the set $\mathbb{Q}_n := \mathbb{Q} \cap I_n$ ($\mathbb{Q}$ is the set of rationals) is dense in βI_n implies that the set $\{M_r, r \in \mathbb{Q}_n\}$ is dense in the subset $\{M_\omega, \omega \in \beta I_n\}$ of $C[0, 1]$. Consequently, putting $\mathbb{Q}_n = \{r_k\}_{k=1}^\infty$ and

$$E_k := \{\omega \in \beta I_n : |M_\omega(t) - M_{r_k}(t)| < 2^{-n} \text{ for all } 0 \leq t \leq 1\}, \quad (6)$$

we have $\beta I_n = \bigcup_{k=1}^\infty E_k$. Now, if $F_m := E_m \setminus (\bigcup_{k=1}^{m-1} E_k)$, $m = 1, 2, \dots$, then F_m are pairwise disjoint and $\beta I_n = \bigcup_{m=1}^\infty F_m$. Define the measure ν_n on σ -algebra of Borel subsets U of the interval I_n by

$$\nu_n(U) := \sum_{\{k: r_k \in U\}} \mu_n(F_k),$$

where μ_n is the probability measure from (4). Since

$$\nu_n(I_n) = \sum_{k=1}^\infty \mu_n(F_k) = \mu_n(\beta I_n) = 1,$$

then ν_n is a probability measure on I_n . Moreover, by (4) and (6), for all $0 \leq t \leq 1$

$$\begin{aligned} \left| t^p - \int_0^{2^{-n}} M_\lambda(t) d\nu_n(\lambda) \right| &= \left| \int_{\beta I_n} M_\omega(t) d\mu_n(\omega) - \int_0^{2^{-n}} M_\lambda(t) d\nu_n(\lambda) \right| \\ &\leq \sum_{k=1}^\infty \left| \int_{F_k} M_\omega(t) d\mu_n(\omega) - \int_{\{r_k\}} M_\lambda(t) d\nu_n(\lambda) \right| \\ &\leq \sum_{k=1}^\infty \left| \int_{\{r_k\}} (M_\lambda(t) + 2^{-n}) d\nu_n(\lambda) - \int_{\{r_k\}} M_\lambda(t) d\nu_n(\lambda) \right| \\ &\leq 2^{-n} \sum_{k=1}^\infty \nu_n(\{r_k\}) = 2^{-n} \nu_n(I_n) = 2^{-n-1}, \end{aligned}$$

and inequality (5) is proved.

Next, for any $s \in (0, 1)$ and $n, j \in \mathbb{N}$ we set

$$a_{j,n} := \int_{s^j 2^{-n}}^{s^{j-1} 2^{-n}} \frac{d\nu_n(\lambda)}{M(\lambda)}. \quad (7)$$

Then, by inequality (5), we have

$$\sum_{j=1}^\infty [a_{j,n}] M(s^j 2^{-n} t) - 2^{-n} < t^p < \sum_{j=1}^\infty [a_{j,n}] M(s^{j-1} 2^{-n} t) + M(t) 2^{-n} / (1-s) + 2^{-n},$$

where by $[z]$ we denote the integer part of a real number z . Choosing now k_n such that

$$\sum_{j=k_n+1}^\infty [a_{j,n}] M(s^{j-1} 2^{-n}) < 2^{-n},$$

as $M(t) \leq M(1) = 1$, we get

$$F_n(st) - 2^{-n+1} < t^p < F_n(t) + 2^{-n} / (1-s) + 2^{-n+1}, \quad 0 \leq t \leq 1, \quad (8)$$

where

$$F_n(t) := \sum_{j=1}^{k_n} [a_{j,n}] M(s^{j-1} 2^{-n} t). \quad (9)$$

Since the right derivative ρ of M (see (3)) is a nondecreasing function and $0 < s < 1$, from (7) it follows that

$$\begin{aligned} F_n(t) - F_n(st) &\leq \sum_{j=1}^{k_n} a_{j,n} (M(s^{j-1}2^{-n}t) - M(s^j2^{-n}t)) \\ &\leq \sum_{j=1}^{k_n} \frac{2^{-n}s^{j-1}(1-s)\rho(s^{j-1}2^{-n})}{M(s^j2^{-n})} \int_{s^j2^{-n}}^{s^{j-1}2^{-n}} d\nu_n(\lambda). \end{aligned}$$

Furthermore, the estimate

$$F(2x) \geq \int_x^{2x} \rho(s) ds \geq x\rho(x), \quad 0 \leq x \leq 1,$$

combined with the hypothesis that M satisfies the Δ_2 -condition at zero, shows that

$$K := \sup_{0 < x \leq 1} \frac{x\rho(x)}{M(x)} < \infty.$$

Hence,

$$F_n(t) - F_n(st) \leq K(1-s) \sum_{j=1}^{k_n} \frac{M(s^{j-1}2^{-n})}{M(s^j2^{-n})} \int_{s^j2^{-n}}^{s^{j-1}2^{-n}} d\nu_n(\lambda).$$

Moreover, one can readily check that the upper Matuszewska-Orlicz index β_M^0 is finite (see also §2.1) and, by its definition, for each $q > \beta_M$ there is a constant $c_0 > 0$ such that

$$M(s^j2^{-n}) \geq c_0 M(s^{j-1}2^{-n})s^q.$$

As a result, since ν_n is a probability measure, we conclude

$$F_n(t) - F_n(st) \leq K(1-s)s^{-q}c_0^{-1} \sum_{j=1}^{k_n} \int_{s^j2^{-n}}^{s^{j-1}2^{-n}} d\nu_n(\lambda) \leq K(1-s)s^{-q}c_0^{-1}. \quad (10)$$

Let $m \in \mathbb{N}$ and $\varepsilon > 0$ be arbitrary. Choose and fix $s \in (0, 1)$ so that

$$K(1-s)s^{-q}c_0^{-1} < \varepsilon/(2m). \quad (11)$$

Then, from (8) and (10) it follows

$$F_n(t) - \frac{\varepsilon}{2m} - 2^{-n+1} < F_n(st) - 2^{-n+1} < t^p, \quad 0 \leq t \leq 1. \quad (12)$$

Now, taking $n \in \mathbb{N}$ satisfying the inequality

$$\frac{2^{-n}}{1-s} + 2^{-n+1} < \frac{\varepsilon}{2m}, \quad (13)$$

from (8) and (12), we obtain

$$F_n(t) - \frac{\varepsilon}{m} < t^p < F_n(t) + \frac{\varepsilon}{m}, \quad 0 \leq t \leq 1. \quad (14)$$

Therefore, for any $c_i \in [0, 1]$, $i = 1, 2, \dots, m$,

$$\sum_{i=1}^m c_i^p - \varepsilon < \sum_{i=1}^m F_n(c_i) < \sum_{i=1}^m c_i^p + \varepsilon,$$

whence for all $c = (c_k)_{k=1}^n \in \mathbb{R}^n$, $c_k \geq 0$,

$$1 - \varepsilon < \sum_{i=1}^m F_n\left(\frac{c_i}{\|c\|_p}\right) < 1 + \varepsilon.$$

Moreover, since F_n is a convex function, from the latter inequality it follows that

$$\sum_{i=1}^m F_n\left(\frac{c_i}{(1+\varepsilon)\|c\|_p}\right) \leq 1$$

and

$$\sum_{i=1}^m F_n\left(\frac{c_i}{(1-\varepsilon)\|c\|_p}\right) > 1.$$

Therefore, by the definition of the norm in an Orlicz sequence space, for every $m \in \mathbb{N}$ and all $c = (c_k)_{k=1}^n \in \mathbb{R}^n$ we have

$$(1 - \varepsilon)\|c\|_p \leq \left\| \sum_{i=1}^m c_i e_i \right\|_{\ell_{F_n}} \leq (1 + \varepsilon)\|c\|_p, \quad (15)$$

where e_i , $i = 1, 2, \dots$, are the canonical unit vectors in ℓ_{F_n} .

Given $m \in \mathbb{N}$ and $\varepsilon > 0$, select s and n to satisfy (11) and (13). For any $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, k_n$ denote by $A_{j,n}^i$ pairwise disjoint subsets of positive integers such that $\text{card } A_{j,n}^i = [a_{j,n}]$. Then, the vectors

$$u_i := 2^{-n} \sum_{j=1}^{k_n} s^{j-1} \sum_{k \in A_{j,n}^i} e_k, \quad i = 1, 2, \dots, m,$$

are copies of an element from l_m . Moreover, by formula (9), we have

$$\left\| \sum_{i=1}^m c_i u_i \right\|_{\ell_M} = \left\| \sum_{i=1}^m c_i e_i \right\|_{\ell_{F_n}}$$

for all $c_i \in \mathbb{R}$. Combining this with (15), we get (1), which completes the proof.

3. Symmetric finite representability of ℓ^p in Orlicz function spaces

Theorem 2

Let M be an Orlicz function satisfying Δ_2 -condition at infinity. Then ℓ^p is symmetrically finitely representable in the Orlicz function space L_M if and only if $p \in [\alpha_M^\infty, \beta_M^\infty]$, i.e., $\mathcal{F}(L_M) = [\alpha_M^\infty, \beta_M^\infty]$.

Proof.

As in the sequence case, we need only to prove the embedding $[\alpha_M^\infty, \beta_M^\infty] \subset \mathcal{F}(L_M)$. More precisely, we have to check that for every $p \in [\alpha_M^\infty, \beta_M^\infty]$, $m \in \mathbb{N}$ and each $\varepsilon > 0$ there exist equimeasurable and disjointly supported functions u_k , $k = 1, 2, \dots, m$, satisfying for all $c = (c_k)_{k=1}^m \in \mathbb{R}^m$ the inequality:

$$(1 + \varepsilon)^{-1}\|c\|_p \leq \left\| \sum_{k=1}^m c_k u_k \right\|_{L_M} \leq (1 + \varepsilon)\|c\|_p \quad (16)$$

First, $t^p \in C_M^\infty \subset C[0, 1]$ and then the same reasoning as in the proof of Theorem 1 shows that and that for every $n \in \mathbb{N}$ there is a probabilistic measure ν_n on $[2^n, \infty)$ such that for all $t \in [0, 1]$

$$\left| t^p - \int_{2^n}^\infty \frac{M(\lambda t)}{M(\lambda)} d\nu_n(\lambda) \right| < 2^{-n}.$$

For any $s > 1$ and $n, j \in \mathbb{N}$ we define

$$a_{j,n} := \int_{s^{j-1}2^n}^{s^j 2^n} \frac{d\mu_n(\lambda)}{M(\lambda)}.$$

Then, by the preceding inequality,

$$\sum_{j=1}^\infty a_{j,n} M(s^{j-1} 2^n t) - 2^{-n} < t^p < \sum_{j=1}^\infty a_{j,n} M(s^j 2^n t) + 2^{-n}.$$

Next, as M satisfies the Δ_2 -condition at infinity, we have

$$M(s^j 2^n t) \leq (1 + 2^{-n}) M(s^{j-1} 2^n t)$$

for all $j \in \mathbb{N}$ and $t \in [0, 1]$ whenever s is sufficiently close to 1. Fixing such a s , we get

$$\sum_{j=1}^\infty a_{j,n} M(s^{j-1} 2^n t) - 2^{-n} < t^p < \sum_{j=1}^\infty (1 + 2^{-n}) a_{j,n} M(s^{j-1} 2^n t) + 2^{-n}.$$

Combining this inequality with the estimate

$$2^{-n} \sum_{j=1}^\infty a_{j,n} M(s^{j-1} 2^n t) < 2^{-2n} + 2^{-n} t^p < 2^{-n+1}, \quad 0 \leq t \leq 1,$$

we deduce

$$\sum_{j=1}^\infty a_{j,n} M(s^{j-1} 2^n t) - 2^{-n} < t^p < \sum_{j=1}^\infty a_{j,n} M(s^{j-1} 2^n t) + 2^{-n+2}. \quad (17)$$

On the other hand, since $M(u) \geq u$ for all $u \geq 1$, we have

$$a_{j,n} \leq \frac{2}{M(2^n s^{j-1})} \leq 2^{-n} s^{-j+1},$$

which implies that

$$\sum_{j=1}^{\infty} a_{j,n} \leq 2^{-n} \sum_{j=1}^{\infty} s^{-j+1} = 2^{-n+1} \cdot \frac{s}{s-1}.$$

Let $m \in \mathbb{N}$ and $\varepsilon > 0$ be arbitrary. Fix n so that

$$\frac{2^{-n+1}s}{s-1} < \frac{1}{m} \quad \text{and} \quad 2^{-n+2}m < \varepsilon. \quad (18)$$

The first of the inequalities (18) allows us to take pairwise disjoint sets $E_j^i \subset [0, 1]$, $j \in \mathbb{N}$, $i = 1, 2, \dots, m$, with $m(E_j^i) = a_{j,n}$. Then, the functions

$$u_i := \sum_{j=1}^{\infty} 2^n s^{j-1} \chi_{E_j^i}$$

are equimeasurable and disjointly supported on $[0, 1]$. Moreover, for all $c_i \in \mathbb{R}$

$$\int_0^1 M\left(\left|\sum_{i=1}^m c_i u_i(t)\right|\right) dt = \sum_{i=1}^m \sum_{j=1}^{\infty} M(2^n s^{j-1} |c_i|) a_{j,n}.$$

Therefore, by (17) and the second inequality in (18), we get

$$\sum_{i=1}^m |c_i|^p - \varepsilon < \int_0^1 M\left(\left|\sum_{i=1}^m c_i u_i(t)\right|\right) dt < \sum_{i=1}^m |c_i|^p + \varepsilon.$$

Repeating further the arguments from the end of the proof of Theorem 1, we come to (16) and so complete the proof.

References

- [1] Tsirel'son B.S. Not every Banach space contains an imbedding of ℓ^p or c_0 . *Functional Analysis and Its Applications*, 1974, vol. 8, no. 2, pp. 138–141. DOI: <https://doi.org/10.1007/BF01078599>. (English; Russian original)
- [2] Krivine J.L. Sous-espaces de dimension finie des espaces de Banach réticulés. *Annals of Mathematics*, 1976, vol. 104, no. 2, pp. 1–29. Available at: https://www.irif.fr/~krivine/articles/Espaces_reticules.pdf.
- [3] Rosenthal H.P. On a theorem of J.L. Krivine concerning block finite representability of ℓ^p in general Banach spaces. *Journal of Functional Analysis*, 1978, vol. 28, pp. 197–225. DOI: [http://dx.doi.org/10.1016/0022-1236\(78\)90086-1](http://dx.doi.org/10.1016/0022-1236(78)90086-1).
- [4] Albiac F., Kalton N.J. Topics in Banach Space Theory. Graduate Texts in Mathematics 233. New York: Springer-Verlag, 2006. 373 p. DOI: <http://dx.doi.org/10.1007/0-387-28142-8>.
- [5] Krein S.G., Petunin Yu.I., Semenov E.M. Interpolation of linear operators. Moscow: Nauka, 1978, 400 p. Available at: <https://elibrary.ru/item.asp?id=21722209>; <https://booksee.org/book/577975>. (In Russ.)
- [6] Lindenstrauss J., Tzafriri L. Classical Banach Spaces, II. Function Spaces. Berlin, Heidelberg, New York: Springer-Verlag, 1979, 243 p. Available at: <https://1lib.education/book/2307307/8b833b?dsource=recommend>.
- [7] Lindenstrauss J., Tzafriri L. Classical Banach Spaces, I. Sequence Spaces. Berlin–New York: Springer-Verlag, 1977. 190 p. Available at: <https://1lib.education/book/2264754/01841c?dsource=recommend>.
- [8] Astashkin S.V. On the finite representability of ℓ^p -spaces in rearrangement invariant spaces. *St. Petersburg Math. J.*, 2012, vol. 23, no. 2, pp. 257–273. DOI: <http://doi.org/10.1090/S1061-0022-2012-01196-9>. (English; Russian original)
- [9] Krasnoselskii M.A., Rutickii Ya.B. Convex functions and Orlicz spaces. Moscow: Gos. izd. fiz.-mat. lit., 1958, 271 p. Available at: <https://1lib.education/book/2078048/983381?id=2078048&secret=983381>. (In Russ.)
- [10] Rao M.M., Ren Z.D. Theory of Orlicz spaces. Monographs and Textbooks in Pure and Applied Mathematics, vol. 146. New York: Marcel Dekker Inc., 1991. 445 p.
- [11] Maligranda L. Orlicz Spaces and Interpolation. Seminars in Mathematics 5. Campinas: University of Campinas, 1989. 206 p.

- [12] Lindenstrauss Y., Tzafriri L. On Orlicz sequence spaces. III. *Israel Journal of Mathematics*, 1973, vol. 14, pp. 368–389. DOI: <https://doi.org/10.1007/BF02771656>.
- [13] Rudin W. *Functional Analysis*. Moscow: Mir, 1975, 443 p. Available at: <https://www.nehudlit.ru/books/funktsionalnyy-analiz.html>.

Научная статья

DOI: 10.18287/2541-7525-2020-26-4-15-24

УДК 517.982.27

Дата: поступления статьи: 14.10.2020
после рецензирования: 16.11.2020
принятия статьи: 25.11.2020

С.В. Асташкин

Самарский национальный исследовательский университет
имени академика С.П. Королева, г. Самара, Российская Федерация
E-mail: astash56@mail.ru. ORCID: <https://orcid.org/0000-0002-8239-5661>

СИММЕТРИЧНАЯ ФИНИТНАЯ ПРЕДСТАВИМОСТЬ ℓ^p В ПРОСТРАНСТВАХ ОРЛИЧА³

АННОТАЦИЯ

Хорошо известно, что банахово пространство может не содержать подпространств, изоморфных хотя бы одному из пространств ℓ^p ($1 \leq p < \infty$) или c^0 (это было показано Цирельсоном в 1974 г.). В то же время по известной теореме Кривина каждое банахово пространство X всегда содержит хотя бы одно из этих пространств *локально*, т. е. существуют конечномерные подпространства в X сколь угодно большой размерности n , изоморфны (равномерно) ℓ_p^n для некоторых $1 \leq p < \infty$ или c_0^n . В этом случае говорят, что ℓ^p (соответственно c^0) *финитно представимо* в X . Основная цель этой статьи — дать характеристику (с полным доказательством) множества тех p , что ℓ^p *симметрично* финитно представимо в любом сепарабельном пространстве Орлича.

Ключевые слова: ℓ^p -пространство; финитная представимость ℓ^p -пространств; симметричная финитная представимость ℓ^p -пространств; функциональное пространство Орлича; пространство последовательностей Орлича; индексы Матушевской — Орлича.

Цитирование. Astashkin S.V. Symmetric finite representability of ℓ^p in Orlicz spaces // Вестник Самарского университета. Естественная серия. 2020. Т. 26, № 4. С. 15–24. DOI: <http://doi.org/10.18287/2541-7525-2020-26-4-15-24>.

Информация о конфликте интересов: автор и рецензенты заявляют об отсутствии конфликта интересов.

© Асташкин С.В., 2020

Асташкин Сергей Владимирович — доктор физико-математических наук, профессор, заведующий кафедрой функционального анализа и теории функций, Самарский национальный исследовательский университет имени академика С.П. Королева, 443086, Российская Федерация, г. Самара, Московское шоссе, 34.

Литература

- [1] Цирельсон Б.С. Не в любое банахово пространство можно вложить ℓ^p или c^0 // Функциональный анализ и его приложения. 1974. Т. 8, №2. С. 57–60. URL: <http://mi.mathnet.ru/faa2331>.
- [2] Krivine J.L. Sous-espaces de dimension finie des espaces de Banach réticulés // *Annals of Mathematics*. 1976. Vol. 104, № 2. P. 1–29. URL: https://www.irif.fr/~krivine/articles/Espaces_reticules.pdf.

³Работа выполнена в рамках внедрения программы развития Научно-образовательного математического центра Приволжского федерального округа, договор № 075-02-2021-1393.

- [3] Rosenthal H.P. On a theorem of J.L. Krivine concerning block finite representability of ℓ^p in general Banach spaces // Journal of Functional Analysis. 1978. Vol. 28. P. 197–225. DOI: [http://dx.doi.org/10.1016/0022-1236\(78\)90086-1](http://dx.doi.org/10.1016/0022-1236(78)90086-1).
- [4] Albiac F., Kalton N.J. Topics in Banach Space Theory. Graduate Texts in Mathematics 233. New York: Springer-Verlag, 2006. 373 p. URL: <http://dx.doi.org/10.1007/0-387-28142-8>.
- [5] Крейн С.Г., Петунин Ю.И., Семенов Е.М. Интерполяция линейных операторов. Москва: Наука, 1978. 400 с. URL: <https://elibrary.ru/item.asp?id=21722209>; <https://booksee.org/book/577975>.
- [6] Lindenstrauss J., Tzafriri L. Classical Banach Spaces, II. Function Spaces. Berlin, Heidelberg, New York: Springer-Verlag, 1979. 243 p. URL: <https://1lib.education/book/2307307/8b833b?dsource=recommend>.
- [7] Lindenstrauss J., Tzafriri L. Classical Banach Spaces, I. Sequence Spaces. Berlin-New York: Springer-Verlag, 1977. 190 p. URL: <https://1lib.education/book/2264754/01841c?dsource=recommend>.
- [8] Асташкин С.В. О финитной представимости ℓ^p -пространств в перестановочно инвариантных пространствах // Алгебра и анализ. 2011. Т. 23, № 2. С. 77–101. URL: <http://mi.mathnet.ru/aa1235>
- [9] Красносельский М.А., Рутницкий Я.Б. Выпуклые функции и пространства Орлича. Москва: Гос. изд-во физ.-мат. лит., 1958. 271 с. URL: <https://1lib.education/book/2078048/983381?id=2078048&secret=983381>.
- [10] Rao M.M., Ren Z.D. Theory of Orlicz spaces. Monographs and Textbooks in Pure and Applied Mathematics, vol. 146 - Marcel Dekker Inc., New York, 1991. 445 p.
- [11] Maligranda L. Orlicz Spaces and Interpolation. Seminars in Mathematics 5. Campinas: University of Campinas, 1989. 206 p.
- [12] Lindenstrauss Y., Tzafriri L. On Orlicz sequence spaces. III // Israel Journal of Mathematics. 1973. Vol. 14. P. 368–389. DOI: <https://doi.org/10.1007/BF02771656>.
- [13] Рудин У. Функциональный анализ. Москва: Мир, 1975. 443 с. URL: <https://www.nehudlit.ru/books/funktsionalnyy-analiz.html>.